

# UAGS Problem Set 1

Wyatt Reeves

**Problem 1** If  $X$  is a subset of  $k[\mathbb{A}^n]$ , define  $V(X)$  to be the subset of  $\mathbb{A}^n$  such that  $p \in V(X)$  if and only if  $f(p) = 0$  for all  $f \in X$ . Prove that if  $I$  is the ideal generated by  $X$ , then  $V(I) = V(X)$ .

**Problem 2** Let  $n = 2$ , and write  $k[\mathbb{A}^2] = k[x, y]$ . Compute and draw:

- $V(x - a, y - b)$
- $V(x^2 - y)$
- $V(x^2 - x)$
- $V(x^2)$

**Problem 3** Let  $A$  be a commutative ring with a multiplicative unit. We say that  $A$  is *Noetherian* if every increasing chain of ideals in  $A$  stabilizes. In other words, for any set of ideals  $\{I_k\}_{k=1}^{\infty}$  such that

$$I_1 \subseteq I_2 \subseteq I_3 \dots$$

we know that there is some  $N$  such that  $I_k = I_N$  for all  $k \geq N$ . Prove that fields are Noetherian. Prove that  $\mathbb{Z}$  is Noetherian. [Hint:  $\mathbb{Z}$  is a Principal Ideal Domain]

**Problem 4** Prove that the following are equivalent:

- $A$  is Noetherian.
- Every collection  $\mathcal{C}$  of ideals of  $A$  has a maximal element.
- Every ideal of  $A$  is finitely generated.

**Problem 5** By filling in the following outline, prove Hilbert's Basis Theorem: if  $A$  is Noetherian, then  $A[x]$  is Noetherian. Conclude that every variety is cut out by finitely many polynomial equations.

**5.a** For any ideal  $I$  in  $A[x]$ , show that the set of leading coefficients of polynomials in  $I$  forms an ideal  $I'$  in  $A$ .

**5.b** Since  $A$  is Noetherian,  $I'$  is finitely generated, say by  $\{a_1 \dots a_n\}$ . Let  $\{p_1 \dots p_n\}$  be polynomials in  $I$  such that the leading coefficient of  $p_i$  is  $a_i$ . Let  $N$  be the largest degree of any  $p_i$ . Show that for any  $f \in I$ , we can write

$$f = \tilde{f} + \sum_{i=1}^n g_i p_i,$$

where  $g_i \in A[x]$  and  $\tilde{f}$  has degree less than  $N$ .

**5.c** Show that for a fixed  $k$ , the set of leading coefficients of polynomials in  $I$  of degree  $k$  forms an ideal  $I_k$  in  $A$ .

**5.d** Let  $\{b_{k,1} \dots b_{k,m_k}\}$  generate  $I_k$ , and let  $q_{k,i}$  be a polynomial in  $I$  with leading coefficient  $b_{k,i}$ . Show that the finite collection of  $p_1 \dots p_n$  and  $q_{k,1} \dots q_{k,m_k}$  for all  $k < N$  generates  $I$ .

**Problem 6 (ZG)** Consider the function  $R : \mathbb{Z}^{>1} \rightarrow \mathbb{N}$  defined by taking  $R(m)$  to be the number of zeros of  $x^2 - 1$  in  $\mathbb{Z}/m\mathbb{Z}$ . Can you give a general formula for computing  $R(m)$ ? [Hint: Chinese Remainder Theorem] More generally, given  $f \in \mathbb{Z}[x]$ , define  $R_f : \mathbb{Z}^{>1} \rightarrow \mathbb{N}$  by  $R_f(m) := \#\{a \in \mathbb{Z}/m\mathbb{Z} : f(a) = 0\}$ . Can you give a general formula for computing  $R_f(m)$ ? What does geometry have to do with all of this?